

Synthesis and Biological Properties of 4-Chloro-N¹-(5-Chloro-1H-Indol-1-yl)-N²-(2-Chlorophenyl)benzene-1,2-Diamine Derivatives

Nikita Shridhar Andhale¹, Rohit Jaysing Bhor^{1,*}, Mahesh Hari Kolhe², Tanishka Anil Pawar¹, Pornima Ramdas Tambe¹, Rupali Mahadev Ghule¹, Sarthak Anil Bhange¹, Bhakti Sharad Thete¹, Pruthviraj Vijay Kute¹, Pratik Vijay Malvade¹, Jyoti Ramkisan Bhagat¹, Poonam Prakash Waykar¹

¹Department of Pharmaceutical Chemistry, Pravara Rural College of Pharmacy Pravaranagar, Rahata, Ahmednagar, Maharashtra, INDIA.

²Department of QA, Pravara Rural College of Pharmacy Pravaranagar, Rahata, Ahmednagar, Maharashtra, INDIA.

ABSTRACT

Background: Five-membered heterocycle with one neighbouring nitrogen atoms is present in the indole nucleus. Heterocyclic substances feature two adjacent benzene and pyrrole ring atoms. Indole are heterocyclic compounds with one oxygen ring that have a benzoannulated - pyrrole ring. Indole derivatives are highly valuable chemical synthesis and medicinal intermediates. Recent years have seen a surge in interest in the effective synthesis of indole blocks and associated functionalization's, particularly the alkylation of indoles. The two following factors will be the main emphasis of this review: i) Recent advancements in the alkylation of indoles, namely those involving the N1-, C2- and C3-positions and the synthesis of indoles, which was facilitated by transition metals. **Materials and Methods:** Benzoic acid and ethanoic acid, propionic acid, hydroxylamine, 2,5 Dichloro aniline are used for the synthesis of indole derivatives were used for the synthesis of indole. **Results:** Compared to indomethacin, the anti-inflammatory results for the substances or their derivatives like N2-(biphenyl-2-yl)-4-chloro-N1-(5-chloro-1H-indol-1-yl)-4-aminobenzene-1,2-diamine NA5 (Scheme 1B); N2-(biphenyl-2-yl)-4-chloro-N1-(5-chloro-1H-indol-1-yl)-3-aminobenzene-1,2-diamine NA6 (Scheme 1B) and N2-(biphenyl-2-yl)-4-chloro-N1-(5-chloro-1H-indol-1-yl)-2-aminobenzene-1,2-diamine NA7 (Scheme 1B). **Conclusion:** Examined were the anti-inflammatory properties of the substances in the title and their derivatives. The compounds containing indole derivatives with an electron-withdrawing group have a higher activity than those containing an electron-donating group, according to research on the link between structure and activity. Summarising the data from the previously reviewed literature, we can say that indole has a broad spectrum of biological activity. Numerous opportunities exist to investigate indole for new therapeutic uses. Researchers worldwide will benefit from this study's coverage of the chemistry of indole derivatives in the design and synthesis of new pharmaceuticals that might be utilised to treat a range of diseases.

Keywords: Benzoic acid, Ethanoic acid, Propionic acid, Hydroxylamine, Egg albumin solution, Phosphate buffered saline.

Correspondence:

Dr. Rohit Jaysing Bhor

Department of Pharmaceutical Chemistry, Pravara Rural College of Pharmacy Pravaranagar, Rahata, Ahmednagar, Maharashtra, INDIA.
Email: rohit.bhor69@gmail.com
ORCID: 0000-0002-7979-3765

Received: 10-04-2024;

Revised: 20-04-2024;

Accepted: 11-06-2024.

INTRODUCTION

Indole, sometimes called benzopyrrole, is an aromatic compound that has ten π -electrons (two from a lone pair on N and eight from π -bonds). On indole, electrophilic substitution happens easily because of high π -electrons delocalization, much like it does on the benzene ring. An essential heterocyclic system, indole supplies the structural framework for alkaloids derived from plants, Strychnine and Lysergic acid Diethylamide (LSD).¹ They

have distinct smells and are colourless and crystalline in nature. When the indole nucleus is added to pharmaceuticals with a physiologically active pharmacophore, it becomes a significant heterocyclic molecule with wide-ranging biological action. As a result, scientists were interested in creating alternative indole scaffolds for the purpose of screening diverse pharmacological actions.² Indole serves as the parent nucleus in a number of natural chemicals, including tryptophan. Higher plants break down tryptophan to create the plant hormone indole-3-acetic acid. Indole derivatives are of great interest. Indoles having a chemical formula of C_8H_7N and a molecular weight of 117.15 g/mol.³ Compared to standard medicine, the molecule with electron-donating and electron-withdrawing groups attached to the indole rings had a stronger inhibitory capacity against



DOI: 10.5530/jyp.2024.16.64

Copyright Information :

Copyright Author (s) 2024 Distributed under Creative Commons CC-BY 4.0

Publishing Partner : EManuscript Tech. [www.emanuscript.in]

bacterial and fungal strains. Indole has been found to possess broad-spectrum antibacterial, anti-inflammatory, analgesic, anti-tubercular, antihypertensive, anticonvulsant and antiviral properties following a thorough examination of several research and review papers.⁴ Indole is manufactured by a range of bacteria and is found in large quantities in the natural world. As a result, various research organisations have worked to find novel antimicrobial medications and the WHO and young scientists worldwide have made significant efforts to treat these infections,⁵ such as anti-inflammatory, antibacterial, antifungal, herbicidal and antiviral activity, however, is represented pharmacologically by indole, along with its derivatives.⁶ In the Figure 1 below, indole was shown. These days, indoles have a broad variety of pharmacological properties. We have attempted to provide a summary of the significant pharmacological activities of derivatives of indole here. The goal of this study is to provide an overview of the current status of indole syntheses, with a special emphasis on the most recent advancements in novel synthetic methodologies.⁷ Preparing natural goods or bioactive chemicals that include the indole unit is emphasised. We give an overview of the synthesis of indole from a historical viewpoint, highlighting the methods used to choose the nitrogen precursors. The review is structured to provide indole synthesis techniques by using various aromatic substrates that include nitrogen-containing functional groups as a source of nitrogen for the indole moiety.⁸ The scant attention paid to some functional groups and typical reactions in earlier evaluations on this subject has led to their being especially emphasised and underlined. The most recent published results are highlighted, although other synthetic methodologies that have been more often utilised and addressed in thorough, outstanding studies on the subject are summarised. We are especially interested in the indolization processes and the various ring closure techniques; we pay little attention to changing the indole structures from the beginning.⁹ Novel indole syntheses are always being found and the scientific community



Figure 1: Structure of Indole derivatives.

places a high value on these discoveries because of the tactical significance of molecules and significant attention and uses because of their biological and pharmacological properties.¹⁰

MATERIALS AND METHODS

Materials

Benzoic acid and ethanoic acid, propionic acid, hydroxylamine, 2,5 Dichloro aniline, Egg albumin solution, Phosphate buffered saline, distilled water/DMSO are used for the synthesis of indole derivatives Every chemical was of the analytical variety. All of the chemicals were bought from Modern Chemicals in Nashik, while others are offered by the college.

Methods

Every derivative of indoles was created using a traditional process. Melting point was confirmed using the capillary open tube technique. The chemicals' purity was examined using Thin-Layer Chromatography (TLC). Using KBr pellets and a Perkin Elmer Spectrum FTIR equipment, IR spectra have been acquired. A Bruker AVANCE III 500 MHz (AV 500) spectrometer was used to record 1 H-NMR spectrum. Physical Data of Indole derivatives were given in Table 1; Scheme of 4-chloro-N¹-(5-chloro-1H-indol-1-yl)-N²-(2-chlorophenyl) benzene-1,2-diamine were given in Figure 2 and Scheme of 4-chloro-N¹-(5-chloro-1H-indol-1-yl)-N²-(2-chlorophenyl) benzene-1,2-diamine derivatives were given in Figure 3.

Experimental Work

Synthesis of 4-chloro-N¹-(5-chloro-1H-indol-1-yl) benzene-1,2-diamine (Scheme 1A)

2 g of 5-chloro-1H-indol-1-amine and 7 mL of 2,5-dichloroaniline were combined in a flask with a round bottom. For 2 hr, the mixture was heated to 100° in a heating mantel. Once the liquid had cooled, 10% sodium hydroxide solution was gradually added until it was barely litmus-alkaline. Reaction flask was rinsed with ice-cold water. After recrystallizing the crude product with cold water, it was cleaned using around 25 mL of cold water.

Synthesis of 4-chloro-N¹-(5-chloro-1H-indol-1-yl)-N²-(2-chlorophenyl)benzene-1,2-diamine: (Scheme 1A)

Add 2 g of the product and two chloroaniline to a round-bottom flask. Heat the reaction mixture for 3 hr at 100°C in a heating mantel. The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated.

Synthesis of 4 N²-(biphenyl-2-yl)-4-chloro-N¹-(5-chloro-1H-indol-1-yl)benzene-1,2-diamine:NA1 (Scheme 1B)

Add benzene to a flask with a round bottom, add 2 g of product and heat the reaction mixture for 3 hr at 100°C in a heating mantel.

The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated.

Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl acetate: NA2 (Scheme 1B)

Add acetic acid to a flask with a round bottom, add 2 g of product and heat the reaction mixture for 3 hr at 100°C in a heating mantel. The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated. Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl propanoate: NA3 (Scheme 1B).

In a round-bottomed flask, 2 g of product and add ethanoic acid and treat the reaction mixture for 4 hr at 100°C in heating mantel. After cooling, the reaction mixture was filtered and recrystallized with ethanol. The Product was washed with ice cold water and filtered and dried in hot air oven for 10 min and later naturally. Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl formate: NA4 (Scheme 1B).

In a round-bottomed flask, 2 g of product and add formic acid and treat the reaction mixture for 3 hr at 100°C in heating mantel. After cooling, the reaction mixture was filtered and recrystallized with ethanol. The Product was washed with ice cold water and filtered and dried in hot air oven for 10 min and later naturally. Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-4-aminobenzene-1,2-diamine:NA5 (Scheme 1B).

In a round-bottomed flask, 2 g of product and add 4 nitro aniline and treat the reaction mixture for 3 hr at 100°C in heating mantel. The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated. Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-3-aminobenzene-1,2-diamine: NA6 (Scheme 1B).

Put 2 g of the product and 3 g of nitro aniline into a round-bottom flask. Heat the reaction mixture for 3 hr at 100°C in a heating mantel. The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated.

Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-2-aminobenzene-1,2-diamine: NA7 (Scheme 1B)

Add 2 g of the product and 2 g of nitro aniline to a round-bottom flask. Heat the reaction mixture for 3 hr at 100°C in a heating mantel. The reaction mixture was cooled, filtered and then recrystallized with ethanol. The product was cleaned in ice-cold

water, filtered and dried for ten minutes in a hot air oven before it naturally dehydrated.

RESULTS

Characterization

Spectral Data

Synthesis of 4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-*N*²-(2-chlorophenyl)benzene-1,2-diamine: (Scheme 1A)

FTIR (KBr) ν cm⁻¹: 1600.92 C=C Stretch (Aromatic), 2952.84 C-H Stretch (Aromatic), 2828.10 C-N bend (Aromatic), 3537.45 N-H Stretch (Aromatic), 2423.21 C-H Stretch (Aliphatic), 771.87 C-Cl Stretch (Aromatic), 1521.84 N-H bend (Aromatic); 1H NMR (400 MHz, DMSO): δ 11.7 N-H (s, 1H, *J*=11.1), δ 12.4 N-H (s, 1H, *J*=11.4), δ 7.3-7.9 Ar C-H (m, 12H, *J*=14.3).

Synthesis of 4-*N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)benzene-1,2-diamine:NA1 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1498.45 C=C Stretch (Aromatic), 1160.01 C-C Stretch (Aromatic), 1301.72 C-N Stretch (Aromatic), 3313 N-H Stretch (Aromatic), 799.5 C-Cl Stretch, 2844 C-H (Aromatic); 1H NMR (400 MHz, DMSO): δ 12.5 N-H (s, 1H, *J*=11.7), 13.3 N-H (s, 1H, *J*=11.4), 7.0-7.8 Ar C-H (m, 17H, *J*=14.3).

Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl acetate: NA2 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1586 C=C Stretch (Aromatic), 1132.01 C-C Stretch (Aromatic), 1332.83 C-N Stretch (Aromatic), 3359 N-H Stretch (Aromatic), 1728.10 C=O (Ester), 782 C-Cl bend (Aliphatic); 1H NMR (400 MHz, DMSO): δ 12.5 N-H (s, 1H, *J*=11.7), 13.3 N-H (s, 1H, *J*=11.4), 7.1-7.9 Ar C-H (m, 15H, *J*=14.3), 2.5 CH₃ (s, 3H, *J*=11.0).

Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl propanoate: NA3 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1718.26 C=C Stretch (Aromatic), 1191.63 C-C Stretch (Aromatic), 1266.86 C-N Stretch (Aromatic), 3259.13 N-H Stretch (Aromatic), 747.0 C-Cl bend (Aliphatic), 1728.15 C=O (Ester); 1H NMR (400 MHz, DMSO): δ 2.2 CH₂ (s, 2H, *J*=11.7), 13.3 N-H (s, 1H, *J*=11.4), 12.7 N-H(s, 1H, *J*=11.4), (7.0-7.8 Ar C-H (m, 13H, *J*=14.3), 2.8 CH₃ (s, 3H, *J*=11.0).

Synthesis of 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl formate: NA4 (Scheme 1B)
FTIR (KBr) ν cm⁻¹: 1632.29 C=C Stretch (Aromatic), 1232.5 R-O-R (Aliphatic), 2835.81 C-N bend (Aromatic), 998.02 N-H Bend (Aromatic), 3247.7 O-H bend (Aliphatic), 1728.1 C=O (Diketones), 1550 N-O (Aromatic), 778.4 C-H bend (Aliphatic); 1H NMR (400 MHz, DMSO): δ 3.3 CH₃ (s, 3H, *J*=11.4), 6.4 CH₂ Group (s, 2H, *J*=13.1), 7.0-7.9 Ar C-H (m, 14H, *J*=14.3); Mol.Wt. 374.35.

Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-4-aminobenzene-1,2-diamine: NA5 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1679.69 C=C Stretch (Aromatic), 1182.78 C-O Stretch (Aromatic), 2828.10 C-N stretch (Aromatic), 1646.80 N-H bend (Aromatic), 3289.2 O-H bend (Aliphatic), 1722.46 C=O (Diketones), 3289.2 O-H Stretch (Aliphatic), 620.66 C-Cl (Aliphatic); 1H NMR (400 MHz, DMSO): δ 2.5 CH₃(s, 3H, *J*=11.7), 3.3 CH₃ (s, 3H, *J*=11.4), 7.7-7.98 Ar C-H (m, 16H, *J*=14.3); Mol.Wt.353.24.

Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-3-aminobenzene-1,2-diamine: NA6 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1594.6 C=C Stretch (Aromatic), 1262.2 C-C Stretch (Aromatic), 2828.10 C-N bend (Aromatic), 1496.6 N-H bend (Aromatic), 3279.2 O-H bend (Aliphatic), 1134.5 C=O (Diketones); 1H NMR (400 MHz, DMSO): δ 2.5 CH₃ (s, 3H, *J*=11.7), 3.3 CH₃ (s, 3H, *J*=11.4), 7.5-7.98 Ar C-H (m, 15H, *J*=14.3); Mol.Wt.374.35.

Synthesis of *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-2-aminobenzene-1,2-diamine: NA7 (Scheme 1B)

FTIR (KBr) ν cm⁻¹: 1495.45 C=C Stretch (Aromatic), 1145.12 C-C Stretch (Aromatic), 2431 C-N Stretch (Aromatic), 3313 N-H Stretch (Aromatic), 1699.5 C=O Stretch (Carboxylic Acid), 3471 and 3537 O-H Stretch (Aliphatic), 1670.10 C=O (Diketones), 2844 C-H(Aromatic); 1H NMR (400 MHz, DMSO): δ 3.63 CH₃ (s, 3H, *J*=11.7), 3.71 CH₃ (s, 3H, *J*=11.4), 5.44 CH₂ Group (s, 2H, *J*=13.1), 6.36-6.38 CH₂ Group (s, 2H, *J*=13.1), 7.18-7.25 Ar C-H (m, 15H, *J*=14.3); 5.98-5.56 OH (2H, s, *J*=17.4); Mol.Wt. 404.46

Biological evaluation**Preparation of 1% of egg albumin solution**

A 1% egg albumin solution may be made using either fresh hen's eggs or easily found egg albumin powder from shops. To correctly make egg-albumin solution from a fresh hen's egg, fracture it gently, then add 1 mL of the transparent part to 100 mL of distilled water with a thorough swirling motion. Egg albumin is the name of the transparent part of the egg. When creating the solution, the water has to be cold. When water reaches a boil, it will coagulate.

Table 1: Physical Data of Indole derivatives

Entry	Yield %	M.P. (°C)	Molecular Formula	Molecular weight
NA	94	130-132	C ₂₀ H ₁₆ N ₂	284.35
NA 1	98	302-304	C ₂₂ H ₂₀ N ₂ O ₂	344.15
NA 2	94	170-173	C ₂₂ H ₂₀ N ₂ O ₄	376.41
NA 3	90	132-134	C ₂₀ H ₁₆ N ₂ O ₂	316.35
NA 4	94	124-126	C ₂₀ H ₁₄ N ₄ O ₂	374.35
NA 5	98	124-126	C ₂₀ H ₁₄ Cl ₂ N ₂	353.24
NA 6	94	184-186	C ₂₀ H ₁₄ N ₄ O ₄	353.24
NA 7	92	166-168	C ₂₄ H ₂₄ N ₂ O ₄	353.24

Table 2: Albumin Denaturation Inhibitory Activity of Indole derivatives (inhibition percentage of protein denaturation).

Indole derivatives	%denaturation inhibition			
	Conc @ 50 μ g/mL	Conc @ 100 μ g/mL	Conc @ 200 μ g/mL	Conc @ 300 μ g/mL
NA	18.25 \pm 1.22	21.12 \pm 1.88	33.22 \pm 2.01	48.13 \pm 3.25
NA 1	19.23 \pm 1.02	24.12 \pm 1.86	34.28 \pm 2.34	50.42 \pm 3.45
NA 2	14.65 \pm 1.24	30.24 \pm 2.01	44.21 \pm 2.96	62.29 \pm 3.25
NA 3	24.36 \pm 1.63	40.27 \pm 2.34	58.22 \pm 3.48	70.35 \pm 3.95
NA 4	23.14 \pm 1.34	36.49 \pm 2.65	51.23 \pm 2.98	56.32 \pm 3.46
NA 5	37.55 \pm 1.96	38.55 \pm 3.54	86.69 \pm 4.08	92.26 \pm 4.66
NA 6	20.25 \pm 1.25	22.25 \pm 1.76	34.22 \pm 2.11	51.13 \pm 3.37
NA 7	21.23 \pm 1.11	21.23 \pm 1.72	36.28 \pm 2.21	52.63 \pm 3.22
Indomethacin		81.53 \pm 3.25	---	---
Control	9.47 \pm 0.04	---	---	---
Negative control (DMSO @ 0.1%)	8.60 \pm 0.03	---	---	---




Figure 2: Scheme of 4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-*N*²-(2-chlorophenyl)benzene-1,2-diamine.

Graphical Representation of Albumin Denaturation Inhibitory Activity of Synthesized Compounds were given in Figure 4.

$$\% \text{denaturation inhibition} = \left[\frac{\text{Abs control} - (\text{Abs test} / \text{Abs control}) \right] \times 100$$

Evaluation of Albumin Denaturation Inhibitory Activity of Synthesized Compounds

Protein denaturation is one of the most well-known causes of inflammation. As part of the investigation, seven test samples (NA1-NA7) had their ability to reduce inflammation assessed. A test for denaturation of proteins has been carried out. Multiple concentrations (50, 100, 200 and 300 µg/mL) of test samples were produced. 0.5 mL of 1.5 mg/mL Bovine Serum Albumin (BSA) was added to each reaction mixture and it was then incubated for 20 min at 37°C. The reaction mixtures were then heated for three minutes to 57°C. Before adding 250 µL of phosphate buffer (0.5 M, pH 6.3), each mixture had to be thoroughly blended. When all of the molecules in each reaction mixture had been evenly distributed, 100 µL of each mixture was transferred into separate test tubes and the same amount of Folin-Ciocalteu's reagent was added. The tubes were allowed to cool after a 10 min incubation time at 55°C. The absorbance at 650 nm was then measured using a UV/vis spectrophotometer, with triple distilled water acting as the blank. The collected data was juxtaposed to the reference medicine, indomethacin sodium (100 µg/mL). The inhibition percentage of protein denaturation was calculated using the procedure given table number 2 below.

DISCUSSION

Research undertaken globally has demonstrated a remarkable influence on scientists who are working on indole derivatives to create indole derivatives that are permitted for use in commerce. Many pharmacological compounds with an indole moiety are being studied for their potential to manage various diseases.¹¹ To establish and support the formulator are the clear features that form the basis and reasoning of the study topic.¹² We reviewed a number of studies published over the past few decades that addressed topics such as anticancer, antifungal, antiplatelet, antidiabetic, antimalarial, antimicrobial, antidiabetic, antiviral, antifungal, anticonvulsant, antifertility, anti-inflammatory, antidepressant, antioxidant, antiestrogenic and antitubercular.¹³⁻¹⁵ Researchers looking for a tactical overview of new indole derivatives with a variety of exciting biological activity and less toxicity and side effects might use this review article as a foundation. Because of its many metabolic activities, indole is an exceptional heterocyclic molecule that is very flexible and has a wide range of pharmacological effects. The amazing indole moiety shares similarities with many different protein architectures.^{16,17} Indole's intriguing molecular structure makes it a good candidate for medication development. Indole derivatives have the ability to bind reversibly to a variety of enzymes and imitate the structure of




Figure 3: Scheme of 4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-*N*²-(2-chlorophenyl)benzene-1,2-diamine derivatives.

peptides, which presents a wealth of options for the development of new medications with unique mechanisms of action. Since the core structures of many compounds contain many physiologically relevant molecules that are active in various illnesses, the existence of nitrogen-based heterocycles in various drugs has been extensively studied.¹⁸ Following scheme 1B, the synthesis of Novel Substituted Indole derivatives from NA1 to NA7 was started. A comparative chart (refer to Table 2 above) demonstrates the anti-inflammatory properties of novel substituted indole-derived compounds at various doses, such as 50, 100, 200 and 300 µg/mL, as well as how well they can stop the spread of inflammation. indomethacin was prescribed as usual medication. Table provided a snapshot of the anti-inflammatory findings for the named compounds (NA1-NA8). Compounds NA5 and NA6 had greater absorbance at lower dosage levels of 50 µg/mL and 100 µg/mL. Compound 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino) phenyl formate NA4 (Scheme 1B) and *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-2-aminobenzene-1,2-diamine NA7 (Scheme 1B) displayed

lower absorbance at higher dose level of 300 µg/mL. Compound 4-*N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)benzene-1,2-diamine NA1 (Scheme 1B) and 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino)phenyl acetate NA2 (Scheme 1B) containing aromatic group, showed higher absorbance at lower dose level of 200 µg/mL. Table displayed the outcomes of the produced compounds' anti-inflammatory tests. The anti-inflammatory properties of recently synthesised compounds have been investigated in relation to the Albumin Denaturation Inhibitory Activity. Tests have been performed on all the substances at 50, 100, 200 and 300 µg/mL. Table 4 summarises the study's findings. It is interesting to point out that all the compounds of the present series showed some anti-inflammatory activity. 2-({5-chloro-2-[(5-chloro-1*H*-indol-1-yl)amino]phenyl}amino) phenyl formate NA4 (Scheme 1B) and *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-2-aminobenzene-1,2-diamine NA7 (Scheme 1B) had more anti-inflammatory




Figure 4: Graphical Representation of Albumin Denaturation Inhibitory Activity of Synthesized Compounds.

properties than the ones that matched them. Nitro aniline compounds showed more activity than other derivatives, while their indole derivatives showed less percentage inhibition by albumin denaturation inhibitory action. It has been observed that the *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-3-aminobenzene-1,2-diamine NA6 (Scheme 1B) and *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-4-aminobenzene-1,2-diamine NA5 (Scheme 1B) and *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-2-aminobenzene-1,2-diamine NA7 (Scheme 1B) having amino at ortho, meta and para position on phenyl ring was the most active and it was studied at three graded doses (50, 100, 200 and 300 µg/mL). The compound *N*²-(biphenyl-2-yl)-4-chloro-*N*¹-(5-chloro-1*H*-indol-1-yl)-4-aminobenzene-1,2-diamine NA5 (Scheme 1B) showed the strongest and most dose-dependent anti-inflammatory action of all the newly created compounds. Because indole derivatives have so many uses in medicinal chemistry, both traditional and green synthetic techniques have been devised for their production. Compared to conventional procedures, the green methodology offers several benefits such as quick reaction times, low costs for reagents, high yields and environmental friendliness. To get beyond the limitations of traditional techniques, researchers in both academia and the pharmaceutical industry are now working on creating a variety of green methodologies for the chemical synthesis of molecules based on indole. This study incorporates the advancements over the past ten years in environmentally friendly techniques for the synthesis of indole derivatives, including the use of microwaves, ionic liquids, water, ultrasound, nanocatalysts, green catalysts, multicomponent reactions and solvent-free reactions.¹⁹ Additionally, this research has demonstrated the use of green chemistry in the production of medicines containing indole and their biological investigations. One of the busiest fields of heterocyclic chemistry research has been indole chemistry. Particularly, 3-substituted indole derivatives have drawn a lot of interest as starting points for the synthesis of several naturally occurring substances as well as additional substances with

biological activity. In this paper, a three-component reaction involving aromatic amine and carboxylic acid with a catalytic quantity of acid under convectional conditions effectively created a synthetic process for multi-substituted indole derivatives. The benefits of this process are high atom utilisation, easy operation and easily accessible raw materials.

CONCLUSION

Convectional technique was used to synthesised several Indole derivatives. The seven Indole derivatives in all were synthesised. Good yields of every chemical were produced. The albumin denaturation inhibitory activity was used to provide the biological anti-inflammatory effect. According to this study, indole derivatives exhibited more potent anti-inflammatory activity against various forms of inflammation. Several of the compounds generated have been demonstrated to have potent anti-inflammatory characteristics. When synthesised compounds were compared to other Indole, higher activity was observed. There are several compounds with a variety of biological applications that possess the indole moiety. Many synthetic medicinal compounds embrace an indole nucleus as part of their pharmacophore structure, which helps pharmaceuticals bind to the residual residues of target molecules' binding sites. Derivatives having indole core have a variety of biological consequences, including antidiabetic, anticancer, antibacterial, anti-HIV, antiviral, anti-inflammatory, antioxidant, anticholinesterase, antitubercular and antimalarial defining features. In their hunt for novel chemical entities, these behaviours have attracted indole to the attention of researchers. These ingredients might be used to create more effective and less hazardous drugs for an assortment of ailments.

ACKNOWLEDGEMENT

The authors are thankful to Dr. S.B. Bhawar, Pravara Rural College of Pharmacy, Pravaranagar.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ABBREVIATIONS

FTIR: Fourier transform infrared spectroscopy; **NMR spectroscopy:** Nuclear magnetic spectroscopy; **MS:** Mass spectroscopy; **KBr:** Potassium Bromide; **%:** Yield Percentage yields; **M.P:** Melting point; **mg/kg:** Milligram/kilograms; **sec:** Seconds; **δ :** Chemical shift; **Mol.Wt:** Molecular Weight gm Gram.

REFERENCES

1. Anger T, Madge DJ, Mulla M, Riddall D. Medicinal chemistry of neuronal voltage-gated sodium channel blockers. *J Med Chem.* 2001;44(2):115-37. doi: 10.1021/jm000155h, PMID 11170622.
2. Ragavendran JV, Sriram D, Kotapati S, Stables J, Yogeewari P. Newer GABA derivatives for the treatment of epilepsy including febrile seizures: A bioisosteric approach. *Eur J Med Chem.* 2008;43(12):2650-55. doi: 10.1016/j.ejmech.2008.01.036, PMID 18403060.
3. Wu T, Ido K, Ohgoh M, Hanada T. Mode of seizure inhibition by sodium channel blockers, an SV2A ligand and an AMPA receptor antagonist in a rat amygdala kindling model. *Epilepsy Res.* 2019;154:42-9. doi: 10.1016/j.eplepsyres.2019.03.011, PMID 31035244.
4. Alexander SM, Harkless J, Butcher JR, Scott KR, Jackson-Ayotunde LP. An Examination of Some Ethyl Ester enaminone Derivatives as Anticonvulsant Agents. *Bioorg Med Chem.* 2013;21:3272-9.
5. Heinbockel T, Wang ZJ, Jackson-Ayotunde PL. Allosteric modulation of GABAA receptors by an anilino enaminone in an olfactory center of the mouse Brain. *Pharmaceuticals (Basel).* 2014;7(12):1069-90. doi: 10.3390/ph7121069, PMID 25525715.
6. Wang ZJ, Sun L, Jackson PL, Scott KR, Heinbockel T. A Substituted anilino enaminone Acts as a Novel Positive allosteric Modulator of GABAA Receptors in the mouse Brain. *J Pharmacol Exp Ther.* 2011;336(3):916-24. doi: 10.1124/jpet.110.173740.
7. Jackson PL, Hanson CD, Farrell AK, Butcher RJ, Stables JP, Eddington ND, *et al.* Enaminones 12. An explanation of anticonvulsant activity and toxicity per Linus Pauling's clathrate hypothesis. *Eur J Med Chem.* 2012;51:42-51. doi: 10.1016/j.ejmech.2012.02.003, PMID 22417639.
8. Anderson AJ, Nicholson JM, Bakare O, Butcher RJ, Wilson TL, Scott KR. Further Studies on the Anticonvulsant Activity and Potential Type IV phosphodiesterase Inhibitory Activity of Substituted Vinylic benzamides. *Bioorg Med Chem.* 2006;14(4):997-1006. doi: 10.1016/j.bmc.2005.09.023, PMID 16219468.
9. Jackson PL, Scott KR, Southerland WM, Fang YY. Enaminones 8: CoMFA and CoMSIA studies on some anticonvulsant enaminones. *Bioorg Med Chem.* 2009;17(1):133-40. doi: 10.1016/j.bmc.2008.11.014, PMID 19059784.
10. Edafigo IO, Kombian SB, Ananthalakshmi KV, Salama NN, Eddington ND, Wilson TL, *et al.* Enaminones: Exploring additional therapeutic activities. *J Pharm Sci.* 2007;96(10):2509-31. doi: 10.1002/jps.20967, PMID 17621683.
11. Amaye IJ, Heinbockel T, Woods J, Wang Z, Martin-Caraballo M, Jackson-Ayotunde P. 6 Hz active anticonvulsant fluorinated N-Benzamide enaminones and their inhibitory neuronal activity. *Int J Environ Res Public Health.* 2018;15(8):1784. doi: 10.3390/ijerph15081784, PMID 30127263.
12. Sula A, Booker J, Ng LC, Naylor CE, DeCaen PG, Wallace BA. The complete structure of an activated open sodium channel. *Nat Commun.* 2017;8:14205. doi: 10.1038/ncomms14205, PMID 28205548.
13. Temme L, Börgel F, Schepmann D, Robaa D, Sippl W, Daniluc C, *et al.* Impact of hydroxy moieties at the benzo annulene ring system of GluN2B ligands: design, synthesis and biological evaluation. *Bioorg Med Chem.* 2019;27(23):115146. doi: 10.1016/j.bmc.2019.115146, PMID 31648876.
14. Yadagiri B, Holagunda UD, Bantu R, Nagarapu L, Guguloth V, Polepally S, *et al.* Rational design, synthesis and anti-proliferative evaluation of novel benzosuberone tethered with hydrazide-hydrazones. *Bioorg Med Chem Lett.* 2014;24(21):5041-4. doi: 10.1016/j.bmcl.2014.09.018, PMID 25264072.
15. Althagafi I, Farghaly TA, Abbas EM, Harras MF. Benzosuberone as precursor for synthesis of antimicrobial agents: synthesis, antimicrobial activity and molecular docking. *Polycyclic Aromat Compd.* 2021;41(8):1646-66. doi: 10.1080/10406638.2019.1692877.
16. Grimaldi TB, Back DF, Zeni G. Potassium tert-butoxide promoted annulation of 2-alkynylphenyl propargyl ethers: selective synthesis of benzofuran and 12 H-benzoannulene derivatives. *J Org Chem.* 2013;78(21):11017-31. doi: 10.1021/jo4020062, PMID 24079836.
17. Ahmed SK, Haese NN, Cowan JT, Pathak V, Moukha-Chafiq O, Smith VJ, *et al.* Targeting Chikungunya virus replication by benzoannulene inhibitors. *J Med Chem.* 2021;64(8):4762-86. doi: 10.1021/acs.jmedchem.0c02183, PMID 33835811.
18. Oh CH, Kim JH, Oh BK, Park JR, Lee JH. Gold-catalyzed sequential activation of propargylic carboxylates: A facile route to Benzoannulenes. *Chemistry.* 2013;19(8):2592-6. doi: 10.1002/chem.201204091, PMID 23296692.
19. Lazzari P, Distinto R, Manca I, Baillie G, Murineddu G, Pira M, *et al.* A critical review of both the synthesis approach and the receptor profile of the 8-chloro-1-(2',4'-dichlorophenyl)-N-piperidin-1-yl-1, 4, 5, 6-tetrahydro benzo [6, 7] cyclohepta [1, 2-c] pyrazole-3-carboxamide and analogue derivatives. *Eur J Med Chem.* 2016;121:194-208. doi: 10.1016/j.ejmech.2016.05.011, PMID 27240274.

Cite this article: Andhale NS, Agale AC, Bhor RJ, Pawar TA, Tambe PR, Ghule RM, *et al.* Synthesis and Biological Properties of 4-Chloro-N1-(5-Chloro-1*H*-Indol-1-yl)-N2-(2-Chlorophenyl) benzene-1,2-Diamine Derivatives. *J Young Pharm.* 2024;16(3):498-505.